Discrete-time ratchets, the Fokker-Planck equation and Parrondo's paradox

PROCEEDINGS THE ROYAL

P. Amengual, A. Allison, R. Toral and D. Abbott

Proc. R. Soc. Lond. A 2004 **460**, 2269-2284 doi: 10.1098/rspa.2004.1283

References

Article cited in: http://rspa.royalsocietypublishing.org/content/460/2048/2269#related-urls

Email alerting service

Receive free email alerts when new articles cite this article - sign up in the box at the top right-hand corner of the article or click $\frac{here}{here}$

MATHEMATICAL, PHYSICAL & ENGINEERING SCIENCES

Discrete-time ratchets, the Fokker–Planck equation and Parrondo's paradox

BY P. Amengual^{1,2}, A. Allison², R. Toral¹ and D. Abbott²

¹Instituto Mediterráneo de Estudios Avanzados, IMEDEA (CSIC-UIB), Ed. Mateu Orfila, Campus UIB, 07122 Palma de Mallorca, Spain (pau@imedea.uib.es)

²Centre for Biomedical Engineering (CBME) and Department of Electrical and Electronic Engineering, The University of Adelaide, SA 5005, Australia

Received 29 August 2003; revised 11 December 2003; accepted 16 January 2004; published online 5 May 2004

Parrondo's games manifest the apparent paradox where losing strategies can be combined to win and have generated significant multidisciplinary interest in the literature. Here we review two recent approaches, based on the Fokker–Planck equation, that rigorously establish the connection between Parrondo's games and a physical model known as the flashing Brownian ratchet. This gives rise to a new set of Parrondo's games, of which the original games are a special case. For the first time, we perform a complete analysis of the new games via a discrete-time Markov chain analysis, producing winning rate equations and an exploration of the parameter space where the paradoxical behaviour occurs.

Keywords: Parrondo's paradox; Fokker-Planck equation; Brownian ratchet

1. Introduction

In many physical and biological systems, combining processes may lead to counterintuitive dynamics. For example, in control theory, the combination of two unstable systems can cause them to become stable (Allison & Abbott 2001*a*). In the theory of granular flow, drift can occur in a counterintuitive direction (Rosato *et al.* 1987; Kestenbaum 1997). Also, the switching between two transient diffusion processes in random media can form a positive recurrent process (Pinsky & Scheutzow 1992). Other interesting phenomena where physical processes drift in a counterintuitive direction can be found (see, for example, Ajdari & Prost 1993; Maslov & Zhang 1998; Westerhoff *et al.* 1986; Key 1987; Abbott 2001).

Parrondo's paradox (Harmer & Abbott 1999*a*, *b*; Harmer *et al.* 2000) is based on the combination of two negatively biased games—losing games—which when combined give rise to a positively biased game, that is, we obtain a winning game. This paradox is a translation of the physical model of the Brownian ratchet into gametheoretic terms. These games were first devised in 1996 by the Spanish physicist Juan M. R. Parrondo, who presented them in unpublished form in Turin, Italy (Parrondo 1996). They served as a pedagogical illustration of the flashing ratchet, where directed motion is obtained from the random or periodic alternation of two relaxation potentials acting on a Brownian particle, neither of which individually produce any net flux (see Reimann (2002) for a complete review on ratchets).

Proc. R. Soc. Lond. A (2004) 460, 2269–2284

 \bigodot 2004 The Royal Society

P. Amengual and others

These games have attracted much interest in other fields, for example, quantum information theory (Abbott *et al.* 2002; Flitney *et al.* 2002; Meyer & Blumer 2002*a*; Lee *et al.* 2002*a*), control theory (Kocarev & Tasev 2002; Dinis & Parrondo 2003), Ising systems (Moraal 2000), pattern formation (Buceta *et al.* 2002*a, b*; Buceta & Lindenberg 2002), stochastic resonance (Allison & Abbott 2001*b*), random walks and diffusions (Cleuren & van den Broeck 2002; Key *et al.* 2002; Kinderlehrer & Kowalczyk 2002; Percus & Percus 2002; Pyke 2002), economics (Boman *et al.* 2001), molecular motors in biology (Ait-Haddou & Herzog 2002; Heath *et al.* 2002) and biogenesis (Davies 2001). They have also been considered as quasi-birth-death processes (Latouche & Taylor 2003) and lattice gas automata (Meyer & Blumer 2002*b*).

Parrondo's two original games are as follows. Game A is a simple coin-tossing game, where a player increases (decreases) his capital in one unit if heads (tails) show up. The probability of winning is denoted by p and the probability of losing is 1 - p.

Game B is a capital-dependent game, where the probability of winning depends upon the actual capital of the player, modulo a given integer M. Therefore, if the capital is i, the probability of winning π_i is taken from the set $\{\pi_0, \pi_1, \ldots, \pi_{M-1}\}$ as $\pi_i = \pi_{i \mod M}$. In the original version of game B, the number M is set equal to three and the probability of winning can take only two values, p_1, p_2 , respectively, according to whether the capital of the player is a multiple of three or not. Using the previous notation we have $p_1 \equiv \pi_0, p_2 \equiv \pi_1 = \pi_2$. The numerical values corresponding to Parrondo's original games (Harmer & Abbott 1999a) are

$$p = \frac{1}{2} - \epsilon,$$

$$p_1 = \frac{1}{10} - \epsilon,$$

$$p_2 = \frac{3}{4} - \epsilon,$$

$$(1.1)$$

where ϵ is a small biasing parameter introduced to control the three probabilities.

Although the original game B was based on a modulo rule, there are other versions of Parrondo's games where this rule has been replaced by a history-dependent rule (Parrondo *et al.* 2000); combinations of two history-dependent games are also considered (Kay & Johnson 2002). Instead of a random alternation, chaotic alternation between the games has been studied (Arena *et al.* 2003). The effects of cooperation between players have also been considered in Parrondo's games, where the probabilities of game B depend on the actual state of the neighbours of the player (Toral 2001). A redistribution of capital between the players has additionally been considered (Toral 2002). Other variations of collective games have recently appeared (Mihailović & Rajković 2003*a*, *b*). For a full review of Parrondo's paradox see Harmer & Abbott (2002).

Games A and B appearing in Parrondo's paradox can be thought of as diffusion processes under the action of an external potential. However, they do not have the more general form of a natural diffusion process, because the capital will always change with every game played, whereas in the most general diffusion process a particle can either move up or down or remain in the same position at a given time. In this article we present a new version of Parrondo's games, where a new transition probability is taken into account. We introduce a *self-transition* probability, that is, the capital of the player now can remain the same after a game played with a probability ρ_i , taken from the set $\{\rho_0, \rho_1, \ldots, \rho_{M-1}\}$ as $\rho_i = \rho_{i \mod M}$. Again, for

simplicity, we will only consider the case of M = 3 with just two possible selftransition probabilities, r_1 , r_2 , depending only on the capital being a multiple of three or not: $r_1 \equiv \rho_0$, $r_2 \equiv \rho_1 = \rho_2$.

As we will show, the significance of this new version is a natural evolution of Parrondo's games, which can now be rigorously derived from the Fokker–Planck equation, based on a physical flashing ratchet model.

The outline of this paper is as follows. In §2 we briefly review two relations concerning Parrondo's games and the Fokker–Planck equation. In both relations the inclusion of self-probabilities is straightforward. In §3 we give a mathematical analysis of the new games using discrete-time Markov chains (DTMCs) and derive conditions for the paradox to appear. In §4 we calculate the rates of winning, describe the parameter space and present numerical simulations which confirm and extend the theoretical analysis. Finally, in §5 we provide a brief discussion of the results.

2. The flashing ratchet and the Fokker–Planck equation

Despite the fact that Parrondo's paradox was inspired by the flashing ratchet, the relation between both has only been made quantitative recently, when two different approaches have established a formal relation between Parrondo's games and the physical model of the flashing ratchet (Allison & Abbott 2002; Toral *et al.* 2003*a*). We now very briefly review both approaches.

In the scheme proposed by Allison & Abbott (2002), the starting point is the following general Fokker–Planck equation (see Horsthemke & Lefever 1984), for the probability P(x,t) of a Brownian particle moving in a time-dependent one-dimensional potential V(x,t):

$$D\frac{\partial^2 P}{\partial x^2} - P\frac{\partial \alpha}{\partial x} - \alpha\frac{\partial P}{\partial x} - \frac{\partial P}{\partial t} = 0, \qquad (2.1)$$

where α and D are referred to as the infinitesimal first and second moments of diffusion, respectively; D has a constant value (Fick's law constant), while $\alpha(x, t)$ is a function related to the applied potential V(x, t) by the equation

$$\alpha(x,t) = -u\frac{\partial}{\partial x}V(x,t), \qquad (2.2)$$

where u denotes the mobility of the Brownian particle.

Equation (2.1) is then discretized using a finite-difference approximation, to obtain

$$P_{i,j} = a_{-1}^{i,j} \cdot P_{i-1,j-1} + a_0^{i,j} \cdot P_{i,j-1} + a_{+1}^{i,j} \cdot P_{i+1,j-1},$$
(2.3)

where

$$a_{-1}^{i,j} = \left\{\frac{D\tau}{\lambda^2} + \frac{\alpha(i,j)\tau}{2\lambda}\right\} \left(\frac{\alpha(i+1,j-1) - \alpha(i-1,j-1)}{2\lambda}\tau + 1\right)^{-1}, \quad (2.4)$$

$$a_0^{i,j} = \left\{ -2\frac{D\tau}{\lambda^2} + 1 \right\} \left(\frac{\alpha(i+1,j-1) - \alpha(i-1,j-1)}{2\lambda} \tau + 1 \right)^{-1},$$
(2.5)

$$a_{+1}^{i,j} = \left\{\frac{D\tau}{\lambda^2} - \frac{\alpha(i,j)\tau}{2\lambda}\right\} \left(\frac{\alpha(i+1,j-1) - \alpha(i-1,j-1)}{2\lambda}\tau + 1\right)^{-1}.$$
 (2.6)

Here the index *i* denotes the discretized space $x = i\lambda$, whereas *j* denotes the discretized time $t = j\tau$; λ and τ account for the space- and time-discretization steps, respectively.

This discretized form (2.3) of the Fokker–Planck equation is compared with the master equation for any of the gambling games used in Parrondo's paradox:

$$P_{i,j} = \pi_{i-1} \cdot P_{i-1,j-1} + \rho_i \cdot P_{i,j-1} + (1 - \pi_{i+1} - \rho_{i+1}) \cdot P_{i+1,j-1}, \qquad (2.7)$$

where $P_{i,j}$ denotes the probability that the player has a capital *i* at the *j*th play. In the original Parrondo games the self-transition probability is zero, so that the term ρ_i is set to zero in the following calculations.

Combining (2.3) and (2.7) we get

$$\frac{\pi_{i-1}}{1-\pi_{i+1}} = \frac{a_{-1}^{i,j}}{a_{+1}^{i,j}} = \left(1 + \frac{\lambda}{2D\tau}\alpha(i,j)\right) \left(1 - \frac{\lambda}{2D\tau}\alpha(i,j)\right)^{-1},$$
(2.8)

and it follows that the function $\alpha(i, j) \equiv \alpha_i$ is independent of the time index j:

$$\alpha_i = \frac{2D}{\lambda} \frac{\pi_{i-1} - (1 - \pi_{i+1})}{\pi_{i-1} + (1 - \pi_{i+1})}.$$
(2.9)

Finally, the discretized values of the potential are obtained by combining (2.2) with (2.9),

$$V_{i} = -\frac{2D}{u} \sum_{k=0}^{i} \left\{ 1 - \left(\frac{1 - \pi_{k+1}}{\pi_{k-1}}\right) \right\} \left(1 + \left(\frac{1 - \pi_{k+1}}{\pi_{k-1}}\right) \right)^{-1}.$$
 (2.10)

This equation allows one to obtain the discretized version of the physical potential V_i in terms of the probabilities π_i of the games.

A second relation between the Fokker–Planck equation and the master equation has been proposed by Toral *et al.* (2003*a*). Unlike the first approach described above, the starting point is now not the Fokker–Planck equation but rather the rewriting of the master equation (2.7) in the form of a continuity equation for the probability:

$$P_{i,j} - P_{i,j-1} = -[J_{i+1,j} - J_{i,j}], \qquad (2.11)$$

where the current $J_{i,j}$ is given by

$$J_{i,j} = \frac{1}{2} [F_i P_{i,j} + F_{i-1} P_{i-1,j}] - [D_i P_{i,j} - D_{i-1} P_{i-1,j}]$$
(2.12)

and

$$F_i = 2\pi_i + \rho_i - 1, \qquad D_i = \frac{1 - \rho_i}{2}.$$
 (2.13)

These coefficients can be related with their analogous terms corresponding to a discretization of the Fokker–Planck equation for a probability P(x,t),

$$\frac{\partial P(x,t)}{\partial t} = -\frac{\partial J(x,t)}{\partial x},\tag{2.14}$$

with a current

$$J(x,t) = F(x)P(x,t) - \frac{\partial [D(x)P(x,t)]}{\partial x}$$
(2.15)

for a general drift F(x) and diffusion D(x).

Again considering the case $\rho_i = 0$, we have

$$D_i \equiv D = \frac{1}{2}, \qquad F_i = -1 + 2\pi_i$$
 (2.16)

and the following form for the current:

$$J_{i,j} = \pi_{i-1} P_{i-1,j} - (1 - \pi_i) P_{i,j}, \qquad (2.17)$$

which is merely the probability flux from state i - 1 to state i.

The relation between the external potential V_i and the games probabilities is in this formulation written as

$$V_i = -\frac{1}{2} \sum_{k=1}^{i} \ln\left[\frac{1+F_{k-1}}{1-F_k}\right] = -\frac{1}{2} \sum_{k=1}^{i} \ln\left[\frac{\pi_{k-1}}{1-\pi_k}\right],$$
(2.18)

where the value $V_0 = 0$ has been adopted for convenience. This equation is the main result concerning the relation between the games probabilities π_i and the discretized version of the potential V_i . As with (2.10), through (2.18) we can obtain the potential that corresponds to a Parrondo game. Notice that both approaches yield different values for the potential V_i corresponding to a set of games probabilities $\{\pi_0, \ldots, \pi_{M-1}\}$. For instance, in the case of a fair game, the potential given by (2.18) is a periodic function $V_{i+M} = V_i$ (Toral *et al.* 2003*b*). Nevertheless, it can be shown that both potentials coincide in the limit of an infinitesimally small space-discretized step λ .

It is possible to solve the master equation (2.11) using a constant current $J_{i,j} = J$, together with the boundary condition $P_i^{\text{st}} = P_{i+M}^{\text{st}}$ in order to obtain the stationary probability distribution P_i^{st} . The result is

$$P_i^{\rm st} = N e^{-2V_i} \left[1 - \frac{2J}{N} \sum_{j=1}^i \frac{e^{2V_j}}{1 - F_j} \right]$$
(2.19)

with a current

$$J = N(e^{-2V_M} - 1) \left(2\sum_{j=1}^{M} \frac{e^{2V_j}}{1 - F_j}\right)^{-1}$$
(2.20)

and N is a normalization constant obtained from

$$\sum_{k=0}^{M-1} P_k^{\mathrm{st}} = 1.$$

The inverse problem of obtaining the game probabilities in terms of the potential can also be solved. It requires the solution of (2.18) with the boundary condition $F_0 = F_M$. The result is given by

$$F_{i} = (-1)^{i} e^{2V_{i}} \left[\frac{\sum_{j=1}^{M} (-1)^{j} [e^{-2V_{j}} - e^{-2V_{j-1}}]}{(-1)^{M} e^{2(V_{0} - V_{M})} - 1} + \sum_{j=1}^{i} (-1)^{j} [e^{-2V_{j}} - e^{-2V_{j-1}}] \right], \quad (2.21)$$

which, via $\pi_i = (1 + F_i)/2$, allows one to obtain the probabilities π_i in terms of the potential V_i . It is clear that the additional condition $\pi_i \in [0, 1] \forall i$ must be fulfilled by any acceptable solution.



Figure 1. Probability trees of the new games A and B. Game A is formed by three branches, denoting the three possibilities of winning (W), remaining in the same state (R) and losing (L). Note that game B has a capital-dependent rule and therefore is not a martingale.

To sum up, we have two approaches, either (2.10) or (2.18), that allow one to obtain the potential corresponding to a set of probabilities $(\pi_0, \ldots, \pi_{M-1})$ defining a Parrondo game. In both approaches it is very easy to introduce self-probabilities $\rho_i \neq 0$. Therefore, we find it interesting to investigate the effect of these terms in Parrondo's paradox. Therefore, we introduce a new branch in the original games (Harmer & Abbott 2002) that accounts for the self-transition probability denoted by r_i . The new diagrams for the games A and B are presented in figure 1. In the next section we will investigate the effect of this new inclusion upon the Parrondo effect.

3. Analysis of the new Parrondo games with self-transitions

(a) Game A

We start with the new game A, where the probability of winning is p, the probability of remaining with the same capital will be denoted as r, and we lose with probability q = 1 - r - p.

Following the same reasoning as Harmer *et al.* (2000), we will calculate the probability f_j that our capital reaches zero in a finite number of plays, supposing that initially we have a given capital of j units. From Markov chain analysis (Karlin 1973) we have that

- (i) $f_j = 1$ for all $j \ge 0$, and so the game is either fair or a losing one, or
- (ii) $f_j < 1$ for all j > 0, in which case the game can be a winning one because there is a certain probability that our capital can grow indefinitely.

We are looking for the set of numbers $\{f_j\}$ that correspond to the minimal nonnegative solution of the equation

$$f_j = p \cdot f_{j+1} + r \cdot f_j + q \cdot f_{j-1}, \quad j \ge 1,$$
 (3.1)

with the boundary condition

$$f_0 = 1.$$
 (3.2)

With a subtle rearrangement, (3.1) can be written in the form

$$f_j = \frac{p}{1-r} \cdot f_{j+1} + \frac{q}{1-r} \cdot f_{j-1}.$$
(3.3)

The solution of (3.3), for the initial condition (3.2), is

$$f_j = A \cdot \left[\left(\frac{1 - p - r}{p} \right)^j - 1 \right] + 1,$$

where A is a constant. For the minimal non-negative solution we obtain

$$f_j = \min\left[1, \left(\frac{1-p-r}{p}\right)^j\right].$$
(3.4)

We can therefore see that the new game A is a winning game for

$$\frac{1-p-r}{p} < 1, \tag{3.5}$$

a losing game for

$$\frac{1-p-r}{p} > 1 \tag{3.6}$$

and a fair game for

$$\frac{1-p-r}{p} = 1.$$
 (3.7)

(b) Game B

We now analyse the new game B. Like game A, we have introduced the probabilities of a self-transition in each state, that is, if the capital is a multiple of three we have a probability r_1 of remaining in the same state, whereas if the capital is not a multiple of three then the probability is r_2 . The rest of the probabilities will follow the same notation as in the original game B, so we have the following scheme:

$$\begin{array}{l} \operatorname{mod}(\operatorname{capital},3) = 0 \to p_1, r_1, q_1, \\ \operatorname{mod}(\operatorname{capital},3) \neq 0 \to p_2, r_2, q_2. \end{array}$$

$$(3.8)$$

As in game A, we will follow similar reasoning as Harmer *et al.* (2000), but this time for game B. Let g_j be the probability that the capital will reach the zeroth state in a finite number of plays, supposing an initial capital of j units. Again, from Markov chain theory we have

- (i) $g_j = 1$ for all $j \ge 0$, so game B is either fair or a losing one, or
- (ii) $g_j < 1$ for all j > 0, in which case game B can be a winning one because there is a certain probability for the capital to grow indefinitely.

The following set of recurrence equations must be solved:

$$g_{3j} = p_1 \cdot g_{3j+1} + r_1 \cdot g_{3j} + (1 - p_1 - r_1) \cdot g_{3j-1}, \qquad j \ge 1, \\ g_{3j+1} = p_2 \cdot g_{3j+2} + r_2 \cdot g_{3j+1} + (1 - p_2 - r_2) \cdot g_{3j}, \qquad j \ge 0, \\ g_{3j+2} = p_2 \cdot g_{3j+3} + r_2 \cdot g_{3j+2} + (1 - p_2 - r_2) \cdot g_{3j+1}, \qquad j \ge 0. \end{cases}$$

$$(3.9)$$

As in game A, we are looking for the set of numbers $\{g_j\}$ that correspond to the minimal non-negative solution. Eliminating terms g_{3j-1} , g_{3j+1} and g_{3j+2} from (3.9) we get

$$[p_1p_2^2 + (1-p_1-r_1)(1-p_2-r_2)^2] \cdot g_{3j} = p_1p_2^2 \cdot g_{3j+3} + (1-p_1-r_1)(1-p_2-r_2)^2 \cdot g_{3j-3}.$$
(3.10)

Considering the same boundary condition as in game A, $g_0 = 1$, the last equation has a general solution of the form

$$g_{3j} = B \cdot \left[\left(\frac{(1 - p_1 - r_1)(1 - p_2 - r_2)^2}{p_1 p_2^2} \right)^j - 1 \right] + 1,$$

where B is a constant. For the minimal non-negative solution we obtain

$$g_{3j} = \min\left[1, \left(\frac{(1-p_1-r_1)(1-p_2-r_2)^2}{p_1p_2^2}\right)^j\right].$$
 (3.11)

It can be verified that the same solution as (3.11) will be obtained by solving (3.9) for g_{3j+1} and g_{3j+2} , leading to the same condition for the probabilities of the games.

As with game A, game B will be a winning one if

(

$$\frac{(1-p_1-r_1)(1-p_2-r_2)^2}{p_1p_2^2} < 1, (3.12)$$

a losing one if

$$\frac{(1-p_1-r_1)(1-p_2-r_2)^2}{p_1p_2^2} > 1,$$
(3.13)

and fair if

$$\frac{(1-p_1-r_1)(1-p_2-r_2)^2}{p_1p_2^2} = 1.$$
(3.14)

(c) Game AB

Now we will turn to the random alternation of games A and B with probability γ . This will be named as game AB. For this game AB we have the following (primed) probabilities:

(i) if the capital is a multiple of three,

$$p_1' = \gamma \cdot p + (1 - \gamma) \cdot p_1,$$

$$r_1' = \gamma \cdot r + (1 - \gamma) \cdot r_1;$$

$$(3.15)$$

(ii) if the capital is not multiple of three,

$$p_2' = \gamma \cdot p + (1 - \gamma) \cdot p_2, r_2' = \gamma \cdot r + (1 - \gamma) \cdot r_2.$$

$$(3.16)$$

The same reasoning as in game B can be made, but using the new probabilities p'_1, r'_1, p'_2, r'_2 instead of p_1, r_1, p_2, r_2 . Eventually, we obtain that game AB will be a winning one if

$$\frac{(1 - p_1' - r_1')(1 - p_2' - r_2')^2}{p_1' p_2'^2} < 1,$$
(3.17)

a losing one if

$$\frac{(1-p_1'-r_1')(1-p_2'-r_2')^2}{p_1'p_2'^2} > 1,$$
(3.18)

and fair if

$$\frac{(1 - p_1' - r_1')(1 - p_2' - r_2')^2}{p_1' p_2'^2} = 1.$$
(3.19)

The paradox will be present if A and B are losing games, while game AB is a winning one. In this framework this means that the conditions (3.6), (3.13) and (3.17) must be satisfied simultaneously. In order to obtain sets of probabilities fulfilling theses conditions, we first obtain sets of probabilities yielding *fair* A and B games but such that AB is a winning game, and then introduce a small biasing parameter ϵ , making A and B losing games, but still keeping a winning AB game. As an example, we give some sets of probabilities that fulfil these conditions:

$$p = \frac{9}{20} - \epsilon, \quad r = \frac{1}{10}, \quad p_1 = \frac{9}{100} - \epsilon, \quad r_1 = \frac{1}{10}, \quad p_2 = \frac{3}{5} - \epsilon, \quad r_2 = \frac{1}{5}, \quad (3.20 \, a)$$

$$p = \frac{9}{20} - \epsilon, \quad r = \frac{1}{10}, \quad p_1 = \frac{509}{5000} - \epsilon, \quad r_1 = \frac{1}{10}, \quad p_2 = \frac{7}{10} - \epsilon, \quad r_2 = \frac{1}{20}, \quad (3.20 \, b)$$

$$p = \frac{9}{20} - \epsilon, \quad r = \frac{1}{10}, \quad p_1 = \frac{3}{25} - \epsilon, \quad r_1 = \frac{2}{10}, \quad p_2 = \frac{3}{5} - \epsilon, \quad r_2 = \frac{1}{10}, \quad (3.20 \, b)$$

$$p = \frac{1}{20} - \epsilon, \quad r = \frac{1}{10}, \quad p_1 = \frac{1}{25} - \epsilon, \quad r_1 = \frac{1}{5}, \quad p_2 = \frac{1}{5} - \epsilon, \quad r_2 = \frac{1}{10}, \quad (3.20\,c)$$

$$p = \frac{1}{4} - \epsilon, \quad r = \frac{1}{2}, \quad p_1 = \frac{3}{25} - \epsilon, \quad r_1 = \frac{2}{5}, \quad p_2 = \frac{3}{5} - \epsilon, \quad r_2 = \frac{1}{10}.$$
(3.20 d)

4. Properties of the games

(a) Rate of winning

If we consider the capital of a player at play number n, X_n modulo M, we can perform a discrete-time Markov chain (DTMC) analysis of the games with a statespace $\{0, 1, \ldots, M - 1\}$ (cf. Harmer *et al.* 2001). For the case of Parrondo's games we have M = 3, so the following set of difference equations for the probability distribution can be obtained (Lee *et al.* 2002b):

$$P_{0,n+1} = p_2 \cdot P_{2,n} + r_1 \cdot P_{0,n} + q_2 \cdot P_{1,n}, P_{1,n+1} = p_1 \cdot P_{0,n} + r_2 \cdot P_{1,n} + q_2 \cdot P_{2,n}, P_{2,n+1} = p_2 \cdot P_{1,n} + r_2 \cdot P_{2,n} + q_1 \cdot P_{0,n},$$

$$(4.1)$$

which can be put in a matrix form as $P_{n+1} = \mathbb{T} \cdot P_n$, where

$$\mathbb{T} = \begin{pmatrix} r_1 & q_2 & p_2 \\ p_1 & r_2 & q_2 \\ q_1 & p_2 & r_2 \end{pmatrix}$$
(4.2)

and

$$\boldsymbol{P}_{n} = \begin{pmatrix} P_{0,n} \\ P_{1,n} \\ P_{2,n} \end{pmatrix}.$$
(4.3)

Proc. R. Soc. Lond. A (2004)

2277

In the limiting case where $n \to \infty$ the system will tend to a stationary state characterized by

$$\boldsymbol{\Pi} = \mathbb{T} \cdot \boldsymbol{\Pi},\tag{4.4}$$

where $\lim_{n\to\infty} \mathbf{P}_n = \boldsymbol{\Pi}$.

Solving (4.4) is equivalent to solving an eigenvalue problem. As we are dealing with Markov chains, we know that there will be an eigenvalue $\lambda = 1$ and the rest will be less than 1 (Karlin 1973). For the $\lambda = 1$ value we obtain the following eigenvector giving the stationary probability distribution in terms of the games' probabilities.

$$\boldsymbol{\Pi} \equiv \begin{pmatrix} \Pi_0 \\ \Pi_1 \\ \Pi_2 \end{pmatrix} = \frac{1}{D} \begin{pmatrix} (1-r_2)^2 - p_2 \cdot (1-p_2-r_2) \\ (1-r_1)(1-r_2) - p_2 \cdot (1-p_1-r_1) \\ (1-r_1)(1-r_2) - p_1 \cdot (1-p_2-r_2) \end{pmatrix},$$
(4.5)

where D is a normalization constant given by

$$D = (1 - r_2)^2 + 2(1 - r_1)(1 - r_2) - p_2(2 - p_2 - r_2 - r_1 - p_1) - p_1(1 - p_2 - r_2).$$
(4.6)

The rate of winning at the nth step has the general expression (Harmer *et al.* 2001)

$$r(n) \equiv E[X_{n+1}] - E[X_n] = \sum_{i=-\infty}^{\infty} i \cdot [P_{i,n+1} - P_{i,n}].$$
(4.7)

Using these expressions and by similar techniques to those employed in Harmer et al. (2001) it is possible to obtain the stationary rate of winning for the new games introduced in the previous section. The results are, for game A,

$$r_{\rm A}^{\rm st} = 2p + r - 1 \tag{4.8}$$

and, for game B,

$$r_{\rm B}^{\rm st} = 2p_2 + r_2 - 1 + [q_2 - p_2 + p_1 - q_1] \cdot \Pi_0$$

= $\frac{3}{D} (p_1 p_2^2 - (1 - p_1 - r_1)(1 - p_2 - r_2)^2),$ (4.9)

where D is given by (4.6).

It is an easy task to check that when $r_1 = r_2 = 0$ we recover the known expressions for the original games obtained by Harmer *et al.* (2001). To obtain the stationary rate for the randomized game AB we just need to replace in the above expression the probabilities from (3.15) and (3.16).

Within this context the paradox appears when $r_{\rm A}^{\rm st} \leq 0$, $r_{\rm B}^{\rm st} \leq 0$ and $r_{\rm AB}^{\rm st} > 0$. If, for example, we use the values from (3.20 *d*) and a switching probability $\gamma = 1/2$, we obtain the following stationary rates for game A, game B and the random combination AB:

$$r_{\rm A}^{\rm st} = -2\epsilon,$$

$$r_{\rm B}^{\rm st} = \frac{-\epsilon(441 - 120\epsilon + 1000\epsilon^2)}{231 - 40\epsilon + 500\epsilon^2},$$

$$r_{\rm AB}^{\rm st} = \frac{93 - 9828\epsilon + 1920\epsilon^2 - 32000\epsilon^3}{2(2499 - 320\epsilon + 8000\epsilon^2)},$$

$$\left.\right\}$$

$$(4.10)$$

which yield the desired paradoxical result for small $\epsilon > 0$.

Proc. R. Soc. Lond. A (2004)

2278

We can also evaluate the stationary rate of winning when both the probability of winning and the self-transition probability for the games vary with a parameter ϵ as $p = p - \frac{1}{2}\epsilon$ and $r = r + \epsilon$, so that normalization is preserved. Using the set of probabilities derived from (3.20 *d*), namely $p = \frac{1}{4} - \frac{1}{2}\epsilon$, $r = \frac{1}{2} + \epsilon$, $p_1 = \frac{3}{25} - \frac{1}{2}\epsilon$, $r_1 = \frac{2}{5} + \epsilon$, $p_2 = \frac{3}{5} - \frac{1}{2}\epsilon$, $r_2 = \frac{1}{10} + \epsilon$, the result is

$$r_{\rm A}^{\rm st} = 0,$$

$$r_{\rm B}^{\rm st} = \frac{-\epsilon(21 - 20\epsilon)}{2(77 - 200\epsilon + 125\epsilon^2)},$$

$$r_{\rm AB}^{\rm st} = \frac{31 - 164\epsilon + 160\epsilon^2}{2(833 - 2600\epsilon + 2000\epsilon^2)},$$

$$(4.11)$$

again a paradoxical result.

A comparison between the expressions for the rates of winning of the original Parrondo games (Harmer *et al.* 2001) and the new games can be made in two ways. The first one consists in comparing two games with the same probabilities of winning, say original game A with probabilities $p = \frac{1}{2}$ and $q = \frac{1}{2}$ and the new game A with probabilities $p_{\text{new}} = \frac{1}{2}$, $r_{\text{new}} = \frac{1}{4}$ and $q_{\text{new}} = \frac{1}{4}$. In this case we can think of the 'old' probability of losing q as taking the place of the *self-transition* probability r_{new} and the new game A than in the original game: remember that the new game A has an extra term r in the rate of winning compared with the original rate, and this extra term is what gives rise to the higher value. The same reasoning applies for game B, leading to the same conclusion.

The other possibility could be to compare the two games with the same probability of losing. In this case, we follow the same reasoning as before, but now we can imagine the 'old' probability of winning as replacing the winning and self-transition probabilities of the new game. What we now obtain is a lower rate of winning for the new game compared with the original one. An easy way of checking this is by rewriting (4.8) and (4.9) as

$$r_{\rm A}^{\rm st} = p - q, r_{\rm B}^{\rm st} = \frac{3}{D} (p_1 p_2^2 - q_1 q_2^2).$$
(4.12)

So for the same value of q but a lower value of p we obtain a lower value for the rates of game A and B.

We now explore the range of probabilities in which the Parrondo effect takes place. We restrict ourselves to the case M = 3 and $\gamma = 1/2$ used in the previous formulae.

The fact that we have introduced three new probabilities complicates the representation of the parameter space as we have six variables altogether, two variables $\{p, r\}$ from game A and four variables $\{p_1, r_1, p_2, r_2\}$ coming from game B. In order to simplify this high number of variables, some probabilities must be set so that a representation in three dimensions will be possible. In our case we will fix the variables $\{r, r_1, r_2\}$ so that the surfaces can be represented in the parameter space $\{p, p_1, p_2\}$.

In figure 2 we can see the resulting region where the paradox exists for the variables $r = \frac{1}{4}$, $r_1 = \frac{1}{8}$ and $r_2 = \frac{1}{10}$. Some animations have shown that the volume where the



Figure 2. Parameter space corresponding to the values $r = \frac{1}{4}$, $r_1 = \frac{1}{8}$ and $r_2 = \frac{1}{10}$. The actual region where the paradox exists is delimited by the plane $p_1 = 0$ and the triangular region situated at the frontal face, where all the planes intersect.



Figure 3. Average gain as a function of the number of games played coming from numerical simulation of Parrondo's games with different sets of probabilities. The notation [a, b] indicates that game A was played a times and game B b times. The gains were averaged over 50 000 realizations of the games. (a) Simulation corresponding to the probabilities (3.20 a) and $\epsilon = \frac{1}{500}$; (b) probabilities (3.20 b) and $\epsilon = \frac{1}{200}$; (c) probabilities (3.20 c) and $\epsilon = \frac{1}{200}$; (d) probabilities (3.20 d) and $\epsilon = \frac{1}{200}$.



Figure 4. Comparison of the theoretical rates of winning (dashed lines) together with the rates obtained through simulations (solid lines). All the simulations were obtained by averaging over 50 000 trials and over all possible initial conditions. (a) The parameters correspond to the ones used in equations (4.10). (b) The parameters correspond to the ones used in equations (4.11).

paradox takes place gradually shrinks to zero as the variables r, r_1 and r_2 increase from zero to their maximum value of unity.

Another interesting fact that we have encountered, which remains an open question, is the impossibility of obtaining the equivalent parameter space to figure 2 with the fixed variables $\{p, p_1, p_2\}$ and with the parameter space variables $\{r, r_1, r_2\}$ instead—it is possible to obtain the planes for games A and B, but not for the randomized game AB.

(b) Simulations and discussion

We have analysed the new games A and B, and obtained the conditions in order to reproduce the Parrondo effect. We now present some simulations to verify that the paradox is present for a different range of probabilities (see figure 3). Some interesting features can be observed from these graphs. First note that the performance of random or deterministic alternation of the games drastically changes with the parameters.

We use the notation [a, b] to indicate that game A was played a times and game B b times. The performance of the deterministic alternations [3, 2] and [2, 2] remain close to one another, as can be seen in figure 3. However the alternation [4, 4] has a low rate of winning because as we play each game four times, that causes the dynamics of games A and B to dominate over the dynamic of alternation, thereby considerably reducing the gain.

The performance of the random alternation is more variable, obtaining in some cases a greater gain than in the deterministic cases (see figure 3c).

In figure 4 a comparison between the theoretical rates of winning for games A, B and AB given by (4.10) and (4.11) and the rates obtained through simulations is presented. It is worth noting the good agreement between both results.

It is also interesting to see how evolves the average gain obtained from the random alternation of game A and game B when varying the mixing parameter γ . In figure 5 we compare both the experimental and theoretical curves. As in the original games, the maximum gain obtained for this set of parameters is obtained for a value around $\gamma \sim \frac{1}{2}$ (Lee *et al.* 2002*b*). For other sets of the game probabilities, though, the optimal γ differs from $\gamma = \frac{1}{2}$.



Figure 5. Comparison between the theoretical and the simulation for the gain vs gamma, for the following set of probabilities: $p = \frac{1}{3}$, $r = \frac{1}{3}$; $p_1 = \frac{3}{25}$, $r_1 = \frac{2}{5}$ and $p_2 = \frac{3}{5}$, $r_2 = \frac{1}{10}$. The simulations were carried out by averaging over 50 000 trials and all possible initial conditions.

5. Conclusion

We have reviewed how the derivation of Parrondo's games from the flashing Brownian ratchet can be rigorously established via the Fokker–Planck equation. This procedure reveals new Parrondo games, of which the original Parrondo games are a special case with self-transitions set to zero. This confirms Parrondo's original intuition based on a flashing ratchet is correct with rigour. We interpreted the self-transitions in terms of particles, in the flashing ratchet, that remain stationary in a given cycle. We then presented a new DTMC analysis for the new games showing that Parrondo's paradox still occurs if the appropriate conditions are fulfilled. New expressions for the rates of winning have been obtained, with the result that within certain conditions a higher rate of winning than in the original games can be obtained. We have also studied how the parameter space where the paradox exists changes with the selftransition variables, and conclude that the parameter space corresponding to the original Parrondo's games is a limiting case of the maximum volume: as the selftransition probabilities increase in value the volume shrinks to zero. However, it is worth noting that despite the volume decreases with increasing the self-transition probabilities, the rates of winning that can be obtained are higher than in the original Parrondo's games.

This work was supported by GTECH Australasia; the Ministerio de Ciencia y Tecnología (Spain) and FEDER, projects BFM2001-0341-C02-01 and BFM2000-1108; P.A. acknowledges support from the Govern Balear, Spain.

References

Abbott, D. 2001 Overview: unsolved problems of noise and fluctuations. *Chaos* 11, 526–538.
Abbott, D., Davies, P. C. W. & Shalizi, C. R. 2002 Order from disorder: the role of noise in creative processes. *Fluct. Noise Lett.* 2, C1–C12.

- Ait-Haddou, R. & Herzog, W. 2002 Force and motion generation of myosin motors: muscle contraction. J. Electromyogr. Kinesiol. 12, 435–445.
- Ajdari, A. & Prost, J. 1992 Drift induced by a periodic potential of low symmetry: pulsed dielectrophoresis. C. R. Acad. Sci. Paris Sér. II 315, 1635–1639.
- Allison, A. & Abbott, D. 2001a Control systems with stochastic feedback. Chaos 11, 715–724.
- Allison, A. & Abbott, D. 2001b Stochastic resonance on a Brownian ratchet. Fluct. Noise Lett. 1, L239–L244.
- Allison, A. & Abbott, D. 2002 The physical basis for Parrondo's games. Fluct. Noise Lett. 2, L327–L342.
- Arena, P., Fazzino, S., Fortuna, L. & Maniscalco, P. 2003 Game theory and non-linear dynamics: the Parrondo paradox case study. *Chaos Solitons Fractals* 17, 545–555.
- Boman, M., Johansson, S. J. & Lyback, D. 2001 Parrondo strategies for artificial traders. In Intelligent agent technology (ed. N. Zhong, J. Liu, S. Ohsuga & J. Bradshaw), pp. 150–159. World Scientific.
- Buceta, J. & Lindenberg, K. 2002 Switching-induced Turing instability. Phys. Rev. E 66, 046202.
- Buceta, J., Lindenberg, K. & Parrondo, J. M. R. 2002a Spatial patterns induced by random switching. *Fluct. Noise Lett.* 2, L21–L29.
- Buceta, J., Lindenberg, K. & Parrondo, J. M. R. 2002b Pattern formation induced by nonequilibrium global alternation of dynamics. *Phys. Rev. E* 66, 036216.
- Cleuren, B. & van den Broeck, C. 2002 Random walks with absolute negative mobility. *Phys. Rev.* E **65**, 030101.
- Davies, P. C. W. 2001 Physics and life: the Abdus Salam memorial lecture. In 6th Trieste Conf. Chemical Evolution, Trieste, Italy (ed. J. Chela-Flores, T. Tobias & F. Raulin), pp. 13–20. Dordrecht: Kluwer Academic.
- Dinis, L. & Parrondo, J. M. R. 2003 Optimal strategies in collective Parondo games. *Europhys. Lett.* 63, 319–325.
- Flitney, A. P., Ng, J. & Abbott, D. 2002 Quantum Parrondo's games. Physica A 314, 35-42.
- Harmer, G. P. & Abbott, D. 1999a Parrondo's paradox. Statist. Sci. 14, 206-213.
- Harmer, G. P. & Abbott, D. 1999b Losing strategies can win by Parrondo's paradox. *Nature* **402**, 864.
- Harmer, G. P. & Abbott, D. 2002 A review of Parrondo's paradox. *Fluct. Noise Lett.* **2**, R71–R107.
- Harmer, G. P., Abbott, D. & Taylor, P. G. 2000 The paradox of Parrondo's games. Proc. R. Soc. Lond. A 456, 247–259.
- Harmer, G. P., Abbott, D., Taylor, P. G. & Parrondo, J. M. R. 2001 Brownian ratchets and Parrondo's games. *Chaos* 11, 705–714.
- Heath, D., Kinderlehrer, D. & Kowalczyk, M. 2002 Discrete and continuous ratchets: from coin toss to molecular motor. *Discrete Contin. Dynam. Syst.* B 2, 153–167.
- Horsthemke, W. & Lefever, R. 1984 Noise-induced transitions: theory and applications in physics, chemistry and biology. Springer.
- Karlin, S. 1973 A first course in stochastic processes. Academic.
- Kay, R. J. & Johnson, N. F. 2002 Winning combinations of history-dependent games. *Phys. Rev.* E 67, 056128.
- Kestenbaum, D. 1997 Sand castles and cocktail nuts. New Scient. 154, 25–28.
- Key, E. S. 1987 Computable examples of the maximal Lyapunov exponent. Probab. Theory Relat. Fields 75, 97–107.
- Key, E. S., Klosek, M. & Abbott, D. 2002 On Parrondo's paradox: how to construct unfair games by composing fair games. (Preprint math/0206151.)
- Kinderlehrer, D. & Kowalczyk, M. 2002 Diffusion-mediated transport and the flashing ratchet. Arch. Ration. Mech. Analysis 161, 149–179.

- Kocarev, L. & Tasev, Z. 2002 Lyapunov exponents, noise-induced synchronization, and Parrondo's paradox. Phys. Rev. E 65, 046215.
- Latouche, G. & Taylor, P. G. 2003 Drift conditions for matrix-analytic models. Math. Operat. Res. 28, 346–360.
- Lee, C. F., Johnson, N. F., Rodriguez, F. & Quiroga, L. 2002a Quantum coherence, correlated noise and Parrondo games. *Fluct. Noise Lett.* 2, L293–L298.
- Lee, Y., Allison, A., Abbott, D. & Stanley, H. E. 2002b Minimal Brownian ratchet: an exactly solvable model. Phys. Rev. Lett. 91, 220601.
- Maslov, S. & Zhang, Y. 1998 Optimal investment strategy for risky assests. Int. J. Theor. Appl. Finance 1, 377–387.
- Meyer, D. A. & Blumer, H. 2002a Quantum Parrondo games: biased and unbiased. Fluct. Noise Lett. 2, L257–L262.
- Meyer, D. A. & Blumer, H. 2002b Parrondo games as a lattice gas automata. J. Stat. Phys. 107, 225–239.
- Mihailović, Z. & Rajković, M. 2003a One dimensional asynchronous cooperative Parrondo's games. Fluct. Noise Lett. 3, L389–L398.
- Mihailović, Z. & Rajković, M. 2003b Synchronous cooperative Parrondo's games. (Preprint condmat/0307212.) (Submitted.)
- Moraal, H. 2000 Counterintuitive behaviour in games based on spin models. J. Phys. A 33, L203–L206.
- Parrondo, J. M. R. 1996 How to cheat a bad mathematician. EEC HC&M Network Workshop on Complexity and Chaos (contract no. ERBCHRX-CT 940546), Turin, Italy.
- Parrondo, J. M. R., Harmer, G. P. & Abbott, D. 2000 New paradoxical games based on Brownian ratchets. *Phys. Rev. Lett.* **85**, 5226–5229.
- Percus, O. E. & Percus, J. K. 2002 Can two wrongs make a right? Coin-tossing games and Parrondo's paradox. Math. Intelligencer 24, 68–72.
- Pinsky, R. & Scheutzow, M. 1992 Some remarks and examples concerning the transient and recurrence of random diffusions. Ann. Inst. H. Poincaré B 28, 519–536.
- Pyke, R. 2002 On random walks related to Parrondo's games. (Preprint math/0206150.)
- Reimann, P. 2002 Brownian motors: noisy transport far from equilibrium. *Phys. Rep.* **361**, 57–265.
- Rosato, A., Strandburg, K. J., Prinz, F. & Swendsen, R. H. 1987 Why the Brazil nuts are on top: size segregation of particulate matter by shaking. *Phys. Rev. Lett.* 58, 1038–1040.
- Toral, R. 2001 Cooperative Parrondo's games. Fluct. Noise Lett. 1, L7–L12.
- Toral, R. 2002 Capital redistribution brings wealth by Parrondo's paradox. Fluct. Noise Lett. 2, L305–L311.
- Toral, R., Amengual, P. & Mangioni, S. 2003*a* Parrondo's games as a discrete ratchet. *Physica* A **327**, 105–110.
- Toral, R., Amengual, P. & Mangioni, S. 2003b A Fokker-Planck description for Parrondo's games. In Proc. SPIE Conf. on Noise in Complex Systems and Stochastic Dynamics, Santa Fe, NM, USA (ed. L. Schimansky-Geier, D. Abbott, A. Neiman & C. van den Broeck), vol. 5114, pp. 309-317. Bellingham, WA: SPIE.
- Westerhoff, H. V., Tsong, T. Y., Chock, P. B., Chen, Y. & Astumian, R. D. 1986 How enzymes can capture and transmit free energy contained in an oscillating electric field. *Proc. Natl Acad. Sci.* 83, 4734–4738.

2284